

Nickname: Clyde Country: Germany
Registration ID: VDH/WCD 5608 Testing date: 2018/2/4

Microchip: 276098106520463 DEU

Breed: Whippet

DNA Identified with standard identification ISAG 2006 markers

Gender: Male

Dog's identity verified from microchip or tattoo by veterinarian or other authorised person during sample taking: Yes

#### Test results - Known disorders in the breed

Disorder	Туре	Mode of Inheritance	Result
Hereditary Phosphofructokinase (PFK) Deficiency	Blood Disorders	Autosomal Recessive	Clear
Myostatin deficiency (Double Muscling, "Bully")	Muscular Disorders	Autosomal Recessive	Clear

### Test results - New potential disorders in the breed

Disorder	Туре	Mode of Inheritance	Result
Factor VII Deficiency	Blood Disorders	Autosomal Recessive	Clear

On behalf of Genoscoper Laboratories,

SIGNATURE





Nickname: Clyde Country: Germany
Registration ID: VDH/WCD 5608 Testing date: 2018/2/4

Microchip: 276098106520463 DEU

Breed: Whippet

DNA Identified with standard identification ISAG 2006 markers

Gender: Male


Dog's identity verified from microchip or tattoo by veterinarian or other authorised person during sample taking: Yes

### Test results for pharmacogenetics

Disorder	Mode of Inheritance	Result
Malignant Hyperthermia (MH)	Autosomal Dominant	Clear
Multi-Drug Resistance 1, (MDR1)	Autosomal Dominant	Clear

On behalf of Genoscoper Laboratories,

SIGNATURE





Nickname: Clyde Country: Germany

Registration ID: VDH/WCD 5608 Testing date: 2018/2/4

Microchip: 276098106520463 DEU

Breed: Whippet

DNA Identified with standard identification ISAG 2006 markers

Gender: Male

Dog's identity verified from microchip or tattoo by veterinarian or other authorised person during sample taking: Yes

### Test results - Traits - page 1

#### **Coat Type**

Trait	Genotype	Description
Coat Length	L/L	The dog is likely to have short-haired coat.
Furnishings / Improper Coat in Portuguese Water Dogs (marker test)	GG/TT	The dog is not genetically likely to express furnishings.
KRT71 c.451C>T (p.Arg151Trp)	C/C	The dog does not carry any copies of the tested allele causing curly coat. The dog most likely has non-curly hair.
MC5R c.237A>T	T/T	The dog has two copies of the allele associated with low shedding. The dog is likely average or low shedder.
SGK3 (p.Val96Glyfs)	1/1	The dog does not carry the tested hairlessness allele of the American Hairless Terrier.

On behalf of Genoscoper Laboratories,

SIGNATURE



Nickname: Clyde Country: Germany

Registration ID: VDH/WCD 5608 Testing date: 2018/2/4

Microchip: 276098106520463 DEU

Breed: Whippet

DNA Identified with standard identification ISAG 2006 markers

Gender: Male

Dog's identity verified from microchip or tattoo by veterinarian or other authorised person during sample taking: Yes

### Test results - Traits - page 2

#### **Coat Colour**

Trait	Genotype	Description
Colour Locus E - Extensions	Em/E	The dog is likely to have a dark mask.
Colour Locus B - Brown	B/B	The dog is not likely to have brown pigment.
Colour Locus K - Dominant Black	KB/ky    KB/kbr    kbr/ky    kbr/kbr	The dog is genetically dominant black or brindle.
Colour Locus A - Agouti	ay/ay	The dog is genetically sable.
Colour Locus S - Piebald or extreme white spotting	sp/sp	The dog is likely to have piebald spotting or to be extreme white.
Colour Locus H - Harlequin	h/h	The dog doesn't have harlequin pattern.
Colour Locus D - Dilution (d <sup>1</sup> allele - marker test available for limited breeds)	D/d	The dog is likely to be non-dilute. The dog carries dilute coat colour.
Merle (M allele)	m/m	The dog is genetically non-merle and does not carry a SILV gene SINE insertion.
Saddle Tan (RALY gene dupl.)	-/-	The dog may have saddle tan pattern if it has also tan point genotype at the A locus.
Albinism (caL-allele)	C/C	The dog does not carry the tested mutation for albinism.

On behalf of Genoscoper Laboratories,

SIGNATURE



Nickname:ClydeCountry:GermanyRegistration ID:VDH/WCD 5608Testing date:2018/2/4

Microchip: 276098106520463 DEU

Breed: Whippet

DNA Identified with standard identification ISAG 2006 markers

Gender: Male

Dog's identity verified from microchip or tattoo by veterinarian or other authorised person during sample taking: Yes

### Test results - Traits - page 3

#### **Body Size**

Trait	Genotype	Description
IGF1 (chr15:41221438)	G/G	The dog is homozygous for the ancestral allele typically associated with large body mass.
IGF1R c.611G>A (p.Arg204His)	G/G	The dog carries two ancestral alleles typically found in larger-sized breeds.
ACSL4 chrX.82919525C>T	C/C	The dog doesn't have the allele associated with large skeletal size and heavy muscling with considerable back fat thickness.
IGSF1 p.Asp768Glu	C/C	The dog doesn't have the allele associated with heavy muscling
FGF4 insertion	D/D	The dog is homozygous for the ancient allele. The dog is likely to have legs of normal length.
STC2 (chr4:39182836)	A/T	The dog carries one copy of the allele associated with reduced body size and one copy of the allele associated with no size reduction.
GHR1 (p.Glu191Lys)	G/G	The dog has two copies of the ancestral allele associated with larger body size.
GHR2 (p.Pro177Leu)	C/C	The dog has two copies of the ancestral allele associated with larger body size.
HMGA2 (chr10:8348804)	G/G	The dog has two copies of the ancestral allele associated with larger body size.

On behalf of Genoscoper Laboratories,

SIGNATURE



Nickname: Clyde Country: Germany

Registration ID: VDH/WCD 5608 Testing date: 2018/2/4

Microchip: 276098106520463 DEU

Breed: Whippet

DNA Identified with standard identification ISAG 2006 markers

Gender: Male

Dog's identity verified from microchip or tattoo by veterinarian or other authorised person during sample taking: Yes

### Test results - Traits - page 4

#### Morphology

Trait	Genotype	Description
BMP3 c.1344C>A (p.Phe448Leu)	C/C	The dog does not carry the tested allele typically associated with shortened head (brachycephaly). The dog is more likely to have an elongated head (dolichocephaly).
chr10:11072007	T/T	The dog does not carry an allele typically associated with floppy ears. The dog is more likely to have pricked than floppy ears.
T c.189C>G (p.lle63Met)	C/C	The dog does not carry the tested bobtail-causing genetic variant. The dog is most likely long-tailed.
EPAS1 (p.Gly305Ser)	G/G	The dog does not carry the tested variant associated with adaptation to high altitudes.
LIMBR1 DC-1	G/G	The dog does not carry the tested allele associated with hind dewclaws in Asian breeds. The dog is not likely to have hind dewclaws.
LIMBR1 DC-2	G/G	The dog does not carry the tested allele associated with hind dewclaws in western breeds. The dog is likely not to have hind dewclaws.

On behalf of Genoscoper Laboratories,

SIGNATURE



### Blood Disorders - page 1

Bleeding disorder due to P2RY12 defect  Canine Cyclic Neutropenia, Cyclic Hematopoiesis, Grey Collie Syndrome, Autosomal Recessive  Clear  Canine Leukocyte Adhesion Deficiency (CLAD), type III  Autosomal Recessive  Clear  Canine Scott Syndrome, (CSS)  Factor IX Deficiency or Hemophilia B; mutation Gly379Glu  Factor IX Deficiency or Hemophilia B; mutation originally found in Airedale Terrier  Factor IX Deficiency or Hemophilia B; mutation originally found in German Wirehaired Pointer  Factor IX Deficiency or Hemophilia B; mutation originally found in German Apso  Factor IX Deficiency or Hemophilia B; mutation originally found in Lhasa Apso  Factor VII Deficiency or Hemophilia A; mutation originally found in Boxer Factor VIII Deficiency or Hemophilia A; mutation originally found in Boxer Factor VIII Deficiency or Hemophilia A; mutation originally found in Boxer Factor VIII Deficiency or Hemophilia A; mutation originally found in German Shepherd Dog Factor VIII Deficiency or Hemophilia A; mutation originally found in Old English Sheepdog Factor VIII Deficiency or Hemophilia A; mutation originally found in Old English Sheepdog Factor VIII Deficiency or Hemophilia A; mutation originally found in Old English Sheepdog Factor VIII Deficiency or Hemophilia A; mutation originally found in Old English Sheepdog Factor VIII Deficiency or Hemophilia A; p.Cys548Tyr mutation originally Factor VIII Deficiency or Hemophilia A; p.Cys548Tyr mutation originally Factor VIII Deficiency or Hemophilia A; p.Cys548Tyr mutation originally Factor VIII Deficiency Autosomal Recessive Clear Factor VII Deficiency Clear Factor VIII Deficiency or Hemophilia A; p.Cys548Tyr mutation originally found in Factor VIII Deficiency Autosomal Recessive Clear Factor VIII Deficiency Clear Factor VIII Deficiency Clear Factor VIII Deficiency Autosomal Recessive Clear Factor VIII Deficiency Clear Factor VIII Deficienc	Disorder	Mode of Inheritance	Result
Conine Leukocyte Adhesion Deficiency (CLAD), type III Autosomal Recessive Clear Canine Scott Syndrome, (CSS) Autosomal Recessive Clear Factor IX Deficiency or Hemophilia B; mutation Gly379Glu X-linked Recessive Clear Factor IX Deficiency or Hemophilia B; mutation originally found in Airedale Terrier Factor IX Deficiency or Hemophilia B; mutation originally found in German X-linked Recessive Clear Wirehaired Pointer Factor IX Deficiency or Hemophilia B; mutation originally found in German X-linked Recessive Wordling Reces	Bleeding disorder due to P2RY12 defect	Autosomal Recessive	Clear
Canine Scott Syndrome, (CSS) Autosomal Recessive Clear Factor IX Deficiency or Hemophilia B; mutation Gly379Glu X-linked Recessive Clear Factor IX Deficiency or Hemophilia B; mutation originally found in Airedale Terrier Factor IX Deficiency or Hemophilia B; mutation originally found in German X-linked Recessive Clear Wirehaired Pointer Factor IX Deficiency or Hemophilia B; mutation originally found in German X-linked Recessive No call Apso Factor IX Deficiency or Hemophilia B; mutation originally found in Lhasa Apso Factor VII Deficiency or Hemophilia A; mutation originally found in Boxer X-linked Recessive Clear Factor VIII Deficiency or Hemophilia A; mutation originally found in Boxer X-linked Recessive Clear Factor VIII Deficiency or Hemophilia A; mutation originally found in X-linked Recessive Clear Factor VIII Deficiency or Hemophilia A; mutation originally found in Old X-linked Recessive Clear Factor VIII Deficiency or Hemophilia A; p.Cys548Tyr mutation originally X-linked Recessive Clear Factor XI Deficiency or Hemophilia A; p.Cys548Tyr mutation originally X-linked Recessive Clear Factor XI Deficiency or Hemophilia A; p.Cys548Tyr mutation originally found in German Shepherd Factor XI Deficiency or Hemophilia A; p.Cys548Tyr mutation originally found in Autosomal Recessive Clear Glanzmann Thrombasthenia Type I, (GT); mutation originally found in Autosomal Recessive Clear Glanzmann Thrombasthenia Type I, (GT); mutation originally found in Autosomal Recessive Clear Macrothrombocytopenia; disease-linked variant originally found in Norfolk and Caim Terrier May-Hegglin Anomaly (MHA) Autosomal Recessive Clear		Autosomal Recessive	Clear
Factor IX Deficiency or Hemophilia B; mutation Gly379Glu X-linked Recessive Clear Factor IX Deficiency or Hemophilia B; mutation originally found in Airedale Terrier Factor IX Deficiency or Hemophilia B; mutation originally found in German Wirehaired Pointer Factor IX Deficiency or Hemophilia B; mutation originally found in Lhasa Apso Factor IX Deficiency or Hemophilia B; mutation originally found in Lhasa Apso Factor VII Deficiency or Hemophilia A; mutation originally found in Boxer Factor VIII Deficiency or Hemophilia A; mutation originally found in Boxer Factor VIII Deficiency or Hemophilia A; mutation originally found in German Shepherd Dog Factor VIII Deficiency or Hemophilia A; mutation originally found in Old English Sheepdog Factor VIII Deficiency or Hemophilia A; mutation originally found in Old English Sheepdog Factor VIII Deficiency or Hemophilia A; p.Cys548Tyr mutation originally Factor VIII Deficiency or Hemophilia A; p.Cys548Tyr mutation originally Factor VIII Deficiency or Hemophilia A; p.Cys548Tyr mutation originally Factor XI Deficiency Autosomal Dominant (Incomplete Penetrance)  Glanzmann Thrombasthenia Type I, (GT); mutation originally found in Pyrenean Mountain Dog  Glanzmann Thrombasthenia Type I, (GT); mutation originally found in Pyrenean Mountain Dog  Glanzmann Thrombasthenia Type I, (GT); mutation originally found in Pyrenean Mountain Type I, (GT); mutation originally found in Pyrenean Mountain Type I, (GT); mutation originally found in Pyrenean Mountain Type I, (GT); mutation originally found in Pyrenean Mountain Type I, (GT); mutation originally found in Pyrenean Mountain Type I, (GT); mutation originally found in Pyrenean Mountain Type I, (GT); mutation originally found in Pyrenean Mountain Type I, (GT); mutation originally found in Pyrenean Mountain Type I, (GT); mutation originally found in Pyrenean Mountain Type I, (GT); mutation originally found in Norfolk Pyrenean Mountain Type I, (GT); mutation originally found in Norfolk Pyrenean Mountain Type I, (GT); mutation Originally foun	Canine Leukocyte Adhesion Deficiency (CLAD), type III	Autosomal Recessive	Clear
Factor IX Deficiency or Hemophilia B; mutation originally found in Airedale X-linked Recessive Clear Terrier  Factor IX Deficiency or Hemophilia B; mutation originally found in German Wirehaired Pointer  Factor IX Deficiency or Hemophilia B; mutation originally found in Lhasa Apso  Factor IXI Deficiency or Hemophilia B; mutation originally found in Lhasa Az-linked Recessive No call Apso  Factor VII Deficiency or Hemophilia A; mutation originally found in Boxer X-linked Recessive Clear  Factor VIII Deficiency or Hemophilia A; mutation originally found in Boxer X-linked Recessive Clear  Factor VIII Deficiency or Hemophilia A; mutation originally found in Serman Shepherd Dog  Factor VIII Deficiency or Hemophilia A; mutation originally found in Old X-linked Recessive Clear  Factor VIII Deficiency or Hemophilia A; p.Cys548Tyr mutation originally X-linked Recessive Clear  Factor XI Deficiency or Hemophilia A; p.Cys548Tyr mutation originally X-linked Recessive Clear  Factor XI Deficiency Or Hemophilia A; p.Cys548Tyr mutation originally X-linked Recessive Clear  Factor XI Deficiency Or Hemophilia A; p.Cys548Tyr mutation originally Autosomal Dominant (Incomplete Penetrance)  Glanzmann Thrombasthenia Type I, (GT); mutation originally found in Autosomal Recessive Clear  Glanzmann Thrombasthenia Type I, (GT); mutation originally found in Autosomal Recessive Clear  Macrothrombocytopenia; disease-linked variant originally found in Norfolk autosomal Recessive Clear  May-Hegglin Anomaly (MHA) Autosomal Dominant Clear  Prekallikrein Deficiency Clear	Canine Scott Syndrome, (CSS)	Autosomal Recessive	Clear
Terrier Factor IX Deficiency or Hemophilia B; mutation originally found in German X-linked Recessive Clear Wirehaired Pointer Factor IX Deficiency or Hemophilia B; mutation originally found in Lhasa Apso Autosomal Recessive Clear Factor VII Deficiency Autosomal Recessive Clear Factor VIII Deficiency or Hemophilia A; mutation originally found in Boxer X-linked Recessive Clear Factor VIII Deficiency or Hemophilia A; mutation originally found in Boxer X-linked Recessive Clear German Shepherd Dog Factor VIII Deficiency or Hemophilia A; mutation originally found in Old X-linked Recessive Clear English Sheepdog Factor VIII Deficiency or Hemophilia A; mutation originally found in Old X-linked Recessive Clear English Sheepdog Factor VIII Deficiency or Hemophilia A; p.Cys548Tyr mutation originally X-linked Recessive Clear Factor XI Deficiency Autosomal Dominant (Incomplete Penetrance) Glanzmann Thrombasthenia Type I, (GT); mutation originally found in Autosomal Recessive Clear Pyrenean Mountain Dog Glanzmann Thrombasthenia Type I, (GT); mutation originally found in Autosomal Recessive Clear Macrothrombocytopenia; disease-linked variant originally found in Norfolk and Caim Terrier May-Hegglin Anomaly (MHA) Autosomal Dominant Clear Prekallikrein Deficiency Autosomal Recessive Clear	Factor IX Deficiency or Hemophilia B; mutation Gly379Glu	X-linked Recessive	Clear
Wirehaired Pointer  Factor IX Deficiency or Hemophilia B; mutation originally found in Lhasa Autosomal Recessive Clear  Factor VII Deficiency Autosomal Recessive Clear  Factor VIII Deficiency or Hemophilia A; mutation originally found in Boxer X-linked Recessive Clear  Factor VIII Deficiency or Hemophilia A; mutation originally found in Sactor VIII Deficiency or Hemophilia A; mutation originally found in Clear German Shepherd Dog  Factor VIII Deficiency or Hemophilia A; mutation originally found in Old English Sheepdog  Factor VIII Deficiency or Hemophilia A; p.Cys548Tyr mutation originally A: Iniked Recessive Clear English Sheepdog  Factor VIII Deficiency or Hemophilia A; p.Cys548Tyr mutation originally X-linked Recessive Clear found in German Shepherd  Factor XI Deficiency Autosomal Dominant (Incomplete Penetrance)  Glanzmann Thrombasthenia Type I, (GT); mutation originally found in Pyrenean Mountain Dog  Glanzmann Thrombasthenia Type I, (GT); mutation originally found in Autosomal Recessive Clear  Macrothrombocytopenia; disease-linked variant originally found in Norfolk Autosomal Recessive Clear  Macrothrombocytopenia; disease-linked variant originally found in Norfolk Autosomal Recessive Clear  May-Hegglin Anomaly (MHA) Autosomal Dominant Clear  Prekallikrein Deficiency Autosomal Recessive Clear		X-linked Recessive	Clear
Apso Factor VII Deficiency Factor VIII Deficiency or Hemophilia A; mutation originally found in Boxer Factor VIII Deficiency or Hemophilia A; mutation originally found in Boxer Factor VIII Deficiency or Hemophilia A; mutation originally found in German Shepherd Dog Factor VIII Deficiency or Hemophilia A; mutation originally found in Old English Sheepdog Factor VIII Deficiency or Hemophilia A; p.Cys548Tyr mutation originally found in German Shepherd  Factor XI Deficiency or Hemophilia A; p.Cys548Tyr mutation originally found in German Shepherd  Factor XI Deficiency Autosomal Dominant (Incomplete Penetrance)  Glanzmann Thrombasthenia Type I, (GT); mutation originally found in Pyrenean Mountain Dog  Glanzmann Thrombasthenia Type I, (GT); mutation originally found in mixed breed dogs  Hereditary Elliptocytosis Clear  Macrothrombocytopenia; disease-linked variant originally found in Norfolk and Cairn Terrier  May-Hegglin Anomaly (MHA) Autosomal Recessive Clear  Prekallikrein Deficiency Autosomal Recessive Clear		X-linked Recessive	Clear
Factor VIII Deficiency or Hemophilia A; mutation originally found in Boxer X-linked Recessive Clear Factor VIII Deficiency or Hemophilia A; mutation originally found in X-linked Recessive Clear Factor VIII Deficiency or Hemophilia A; mutation originally found in Old English Sheepdog Factor VIII Deficiency or Hemophilia A; p.Cys548Tyr mutation originally X-linked Recessive Clear Factor VIII Deficiency or Hemophilia A; p.Cys548Tyr mutation originally X-linked Recessive Clear Factor XI Deficiency Autosomal Dominant (Incomplete Penetrance) Glanzmann Thrombasthenia Type I, (GT); mutation originally found in Pyrenean Mountain Dog Glanzmann Thrombasthenia Type I, (GT); mutation originally found in Autosomal Recessive Clear Macrothrombocytopenia; disease-linked variant originally found in Norfolk and Caim Terrier May-Hegglin Anomaly (MHA) Autosomal Dominant Clear Prekallikrein Deficiency Autosomal Recessive Clear		X-linked Recessive	No call
Factor VIII Deficiency or Hemophilia A; mutation originally found in Schepherd Dog  Factor VIII Deficiency or Hemophilia A; mutation originally found in Old English Sheepdog  Factor VIII Deficiency or Hemophilia A; p.Cys548Tyr mutation originally found in Old English Sheepdog  Factor VIII Deficiency or Hemophilia A; p.Cys548Tyr mutation originally found in German Shepherd  Factor XI Deficiency  Autosomal Dominant (Incomplete Penetrance)  Glanzmann Thrombasthenia Type I, (GT); mutation originally found in Pyrenean Mountain Dog  Glanzmann Thrombasthenia Type I, (GT); mutation originally found in mixed breed dogs  Hereditary Elliptocytosis  Clear  Macrothrombocytopenia; disease-linked variant originally found in Norfolk and Cairn Terrier  May-Hegglin Anomaly (MHA)  Autosomal Recessive  Clear  Prekallikrein Deficiency  Autosomal Recessive  Clear	Factor VII Deficiency	Autosomal Recessive	Clear
Factor VIII Deficiency or Hemophilia A; mutation originally found in Old English Sheepdog  Factor VIII Deficiency or Hemophilia A; p.Cys548Tyr mutation originally Tactor VIII Deficiency or Hemophilia A; p.Cys548Tyr mutation originally Tactor VIII Deficiency or Hemophilia A; p.Cys548Tyr mutation originally Tactor VIII Deficiency Clear Sector XI Deficiency Autosomal Dominant (Incomplete Penetrance)  Glanzmann Thrombasthenia Type I, (GT); mutation originally found in Pyrenean Mountain Dog  Glanzmann Thrombasthenia Type I, (GT); mutation originally found in Autosomal Recessive Clear mixed breed dogs  Hereditary Elliptocytosis Clear  Macrothrombocytopenia; disease-linked variant originally found in Norfolk autosomal Recessive Clear  May-Hegglin Anomaly (MHA) Autosomal Dominant Clear Prekallikrein Deficiency Autosomal Recessive Clear	Factor VIII Deficiency or Hemophilia A; mutation originally found in Boxer	X-linked Recessive	Clear
English Sheepdog  Factor VIII Deficiency or Hemophilia A; p.Cys548Tyr mutation originally found in German Shepherd  Factor XI Deficiency  Autosomal Dominant (Incomplete Penetrance)  Glanzmann Thrombasthenia Type I, (GT); mutation originally found in Pyrenean Mountain Dog  Glanzmann Thrombasthenia Type I, (GT); mutation originally found in Autosomal Recessive  Glanzmann Thrombasthenia Type I, (GT); mutation originally found in Autosomal Recessive  Clear Macrothrombocytopenia; disease-linked variant originally found in Norfolk and Cairn Terrier  May-Hegglin Anomaly (MHA)  Autosomal Recessive  Clear  Autosomal Recessive  Clear  Autosomal Recessive  Clear		X-linked Recessive	Clear
Factor XI Deficiency  Autosomal Dominant (Incomplete Penetrance)  Glanzmann Thrombasthenia Type I, (GT); mutation originally found in Pyrenean Mountain Dog  Glanzmann Thrombasthenia Type I, (GT); mutation originally found in Autosomal Recessive  Glear Preeditary Elliptocytosis  Hereditary Elliptocytosis  Clear  Macrothrombocytopenia; disease-linked variant originally found in Norfolk and Caim Terrier  May-Hegglin Anomaly (MHA)  Autosomal Recessive  Clear  Autosomal Dominant  Clear  Prekallikrein Deficiency  Autosomal Recessive  Clear		X-linked Recessive	Clear
Glanzmann Thrombasthenia Type I, (GT); mutation originally found in Pyrenean Mountain Dog  Glanzmann Thrombasthenia Type I, (GT); mutation originally found in Mutosomal Recessive  Glanzmann Thrombasthenia Type I, (GT); mutation originally found in Mutosomal Recessive  Clear  Hereditary Elliptocytosis  Clear  Macrothrombocytopenia; disease-linked variant originally found in Norfolk and Cairn Terrier  May-Hegglin Anomaly (MHA)  Autosomal Recessive  Clear  Prekallikrein Deficiency  Autosomal Recessive  Clear		X-linked Recessive	Clear
Pyrenean Mountain Dog  Glanzmann Thrombasthenia Type I, (GT); mutation originally found in mixed breed dogs  Hereditary Elliptocytosis  Clear  Macrothrombocytopenia; disease-linked variant originally found in Norfolk and Cairn Terrier  May-Hegglin Anomaly (MHA)  Autosomal Dominant  Clear  Prekallikrein Deficiency  Autosomal Recessive  Clear	Factor XI Deficiency		Clear
mixed breed dogs  Hereditary Elliptocytosis  Clear  Macrothrombocytopenia; disease-linked variant originally found in Norfolk and Cairn Terrier  May-Hegglin Anomaly (MHA)  Autosomal Dominant  Clear  Prekallikrein Deficiency  Autosomal Recessive  Clear		Autosomal Recessive	Clear
Macrothrombocytopenia; disease-linked variant originally found in Norfolk and Cairn Terrier  May-Hegglin Anomaly (MHA)  Autosomal Dominant  Clear  Prekallikrein Deficiency  Autosomal Recessive  Clear		Autosomal Recessive	Clear
and Cairn Terrier  May-Hegglin Anomaly (MHA)  Autosomal Dominant  Clear  Prekallikrein Deficiency  Autosomal Recessive  Clear	Hereditary Elliptocytosis		Clear
Prekallikrein Deficiency Autosomal Recessive Clear		Autosomal Recessive	Clear
•	May-Hegglin Anomaly (MHA)	Autosomal Dominant	Clear
Pyruvate Kinase Deficiency; mutation originally found in Basenji Autosomal Recessive Clear	Prekallikrein Deficiency	Autosomal Recessive	Clear
	Pyruvate Kinase Deficiency; mutation originally found in Basenji	Autosomal Recessive	Clear



### Blood Disorders - page 2

Disorder	Mode of Inheritance	Result
Pyruvate Kinase Deficiency; mutation originally found in Beagle	Autosomal Recessive	Clear
Pyruvate Kinase Deficiency; mutation originally found in Pug	Autosomal Recessive	Clear
Pyruvate Kinase Deficiency; mutation originally found in West Highland White Terrier	Autosomal Recessive	Clear
Thrombopathia; mutation originally found in Basset Hound	Autosomal Recessive	Clear
Thrombopathia; mutation originally found in Eskimo Spitz	Autosomal Recessive	Clear
Thrombopathia; mutation originally found in Landseer	Autosomal Recessive	Clear
Trapped Neutrophil Syndrome, (TNS)	Autosomal Recessive	Clear
Von Willebrand's Disease (vWD) Type 1	Autosomal Recessive	Clear
Von Willebrand's Disease (vWD) Type 3; mutation originally found in Kooikerhondje	Autosomal Recessive	Clear
Von Willebrand's Disease (vWD) Type 3; mutation originally found in Scottish Terrier	Autosomal Recessive	Clear
Von Willebrand's Disease (vWD) Type 3; mutation originally found in Shetland Sheepdog	Autosomal Recessive	Clear



### Ocular Disorders - page 1

Disorder	Mode of Inheritance	Result
Canine Multifocal Retinopathy 1, (CMR1); mutation originally found in Mastiff-related breeds	Autosomal Recessive	Clear
Canine Multifocal Retinopathy 2, (CMR2); mutation originally found in Coton de Tulear	Autosomal Recessive	Clear
Canine Multifocal Retinopathy 3, (CMR3); mutation originally found in Lapponian Herder	Autosomal Recessive	Clear
Cone Degeneration, (CD) or Achromatopsia; mutation originally found in Alaskan Malamute	Autosomal Recessive	Clear
Cone Degeneration, (CD) or Achromatopsia; mutation originally found in German Shepherd Dog	Autosomal Recessive	Clear
Cone Degeneration, (CD) or Achromatopsia; mutation originally found in German Shorthaired Pointer	Autosomal Recessive	Clear
Cone-Rod Dystrophy 1, (crd1); mutation originally found in American Staffordshire Terrier	Autosomal Recessive	Clear
Cone-Rod Dystrophy 2, (crd2); mutation originally found in American Pit Bull Terrier	Autosomal Recessive	Clear
Cone-Rod Dystrophy, (cord1-PRA / crd4)	Autosomal Recessive (Incomplete Penetrance)	Clear
Cone-Rod Dystrophy, Standard Wirehaired Dachshund, (crd SWD)	Autosomal Recessive	Clear
Congenital Eye Disease; mutation originally found in Irish Soft-Coated Wheaten Terrier	Autosomal Recessive	Clear
Dominant Progressive Retinal Atrophy, (DPRA)	Autosomal Dominant	Clear
Generalized Progressive Retinal Atrophy	Autosomal Recessive	Clear
Golden Retriever Progressive Retinal Atrophy 1, (GR_PRA 1)	Autosomal Recessive	Clear
Primary Hereditary Cataract, (PHC); mutation originally found in Australian Shepherd	Autosomal Dominant (Incomplete Penetrance)	Clear
Primary Lens Luxation, (PLL)	Autosomal Recessive	Clear
Primary Open Angle Glaucoma, (POAG); mutation originally found in Basset Fauve de Bretagne	Autosomal Recessive	Clear
Primary Open Angle Glaucoma, (POAG); mutation originally found in Beagle	Autosomal Recessive	Clear
Primary Open Angle Glaucoma, (POAG); mutation originally found in Norwegian Elkhound	Autosomal Recessive	Clear
Primary Open Angle Glaucoma, (POAG); mutation originally found in Petit Basset Griffon Vendeen	Autosomal Recessive	Clear
Primary lens luxation (PLL) and glaucoma; mutation originally found in Shar Pei	Autosomal Recessive	Clear



### Ocular Disorders - page 2

Disorder	Mode of Inheritance	Result
Progressive Retinal Atrophy Type III, (PRA type III); mutation originally found in Tibetan Spaniel and Tibetan Terrier	Autosomal Recessive	Clear
Progressive Retinal Atrophy, (CNGA1-PRA); mutation originally found in Shetland Sheepdog	Autosomal Recessive	Clear
Progressive Retinal Atrophy, (PAP1_PRA); mutation originally found in Papillon and Phalene	Autosomal Recessive	Clear
Progressive Retinal Atrophy, (PRA); mutation originally found in Basenji	Autosomal Recessive	Clear
Progressive Retinal Atrophy, (PRA); mutation originally found in Swedish Vallhund	Autosomal Recessive	Clear
Rod-Cone Dysplasia 1, (rcd1); mutation originally found in Irish Setter	Autosomal Recessive	Clear
Rod-Cone Dysplasia 1a, (rdc1a); mutation originally found in Sloughi	Autosomal Recessive	Clear
Rod-Cone Dysplasia 3, (rcd3)	Autosomal Recessive	Clear
X-Linked Progressive Retinal Atrophy 1, (XLPRA1)	X-linked Recessive	Clear
X-Linked Progressive Retinal Atrophy 2, (XLPRA2; Type A PRA)	X-linked Recessive	Clear

#### **Cardiac Disorders**

Disorder	Mode of Inheritance	Result
Dilated Cardiomyopathy, (DCM); mutation originally found in Schnauzer	Autosomal Recessive	Clear
Long QT Syndrome	Autosomal Dominant	Clear

#### **Endocrine Disorders**

Disorder	Mode of Inheritance	Result
Congenital Hypothyroidism; mutation originally found in Tenterfield Terrier	Autosomal Recessive	Clear
Congenital Hypothyroidism; mutation originally found in Toy Fox and Rat Terrier	Autosomal Recessive	Clear



### Immunological Disorders

Disorder	Mode of Inheritance	Result
Autosomal Recessive Severe Combined Immunodeficiency, (ARSCID)	Autosomal Recessive	Clear
Complement 3 (C3) Deficiency	Autosomal Recessive	Clear
Myeloperoxidase Deficiency	Autosomal Recessive	Clear
Severe Combined Immunodeficiency in Frisian Water Dogs, (SCID)	Autosomal Recessive	Clear
X-Linked Severe Combined Immunodeficiency (XSCID); mutation originally found in Basset Hound	X-linked Recessive	Clear
X-Linked Severe Combined Immunodeficiency (XSCID); mutation originally found in Cardigan Welsh Corgi	X-linked Recessive	Clear



#### **Renal Disorders**

Disorder	Mode of Inheritance	Result
Cystinuria Type I-A; mutation originally found in Newfoundland Dog	Autosomal Recessive	Clear
Cystinuria Type II-A; mutation originally found in Australian Cattle Dog	Autosomal Dominant	Clear
Cystinuria, Type II-B; mutation originally found in Miniature Pinscher	Autosomal Dominant	Clear
Fanconi Syndrome	Autosomal Recessive	Clear
Hyperuricosuria, (HUU)	Autosomal Recessive	Clear
Polycystic Kidney Disease in Bull Terriers, (BTPKD)	Autosomal Dominant	Clear
Primary Hyperoxaluria, (PH); mutation originally found in Coton de Tulear	Autosomal Recessive	Clear
Protein Losing Nephropathy, (PLN); NPHS1 gene variant		Clear
Renal Cystadenocarcinoma and Nodular Dermatofibrosis, (RCND)	Autosomal Dominant	Clear
X-Linked Hereditary Nephropathy, (XLHN)	X-linked Recessive	Clear
X-Linked Hereditary Nephropathy, (XLHN); mutation originally found in Navasota Dog	X-linked Recessive	Clear
Xanthinuria, Type 1a; mutation originally found in mixed breed dogs	Autosomal Recessive	Clear
Xanthinuria, Type 2a; mutation originally found in Toy Manchester Terrier	Autosomal Recessive	Clear
Xanthinuria, Type 2b; mutation originally found in Cavalier King Charles Spaniel and English Cocker Spaniel	Autosomal Recessive	Clear



#### **Metabolic Disorders**

Disorder	Mode of Inheritance	Result
Glycogen Storage Disease Type II or Pompe's Disease, (GSD II)	Autosomal Recessive	Clear
Glycogen Storage Disease Type IIIa, (GSD IIIa)	Autosomal Recessive	Clear
Glycogen Storage Disease Type Ia, (GSD Ia)	Autosomal Recessive	Clear
Hypocatalasia or Acatalasemia	Autosomal Recessive	Clear
Intestinal Cobalamin Malabsorption or Imerslund-Gräsbeck Syndrome, (IGS); mutation originally found in Beagle	Autosomal Recessive	Clear
Intestinal Cobalamin Malabsorption or Imerslund-Gräsbeck Syndrome, (IGS); mutation originally found in Border Collie	Autosomal Recessive	Clear
Mucopolysaccharidosis Type IIIA, (MPS IIIA); mutation originally found in Dachshund	Autosomal Recessive	Clear
Mucopolysaccharidosis Type IIIA, (MPS IIIA); mutation originally found in New Zealand Huntaway	Autosomal Recessive	Clear
Mucopolysaccharidosis Type VII, (MPS VII); mutation originally found in Brazilian Terrier	Autosomal Recessive	Clear
Mucopolysaccharidosis Type VII, (MPS VII); mutation originally found in German Shepherd	Autosomal Recessive	Clear
Pyruvate Dehydrogenase Phosphatase 1 (PDP1) Deficiency	Autosomal Recessive	Clear



#### **Muscular Disorders**

Disorder	Mode of Inheritance	Result
Cavalier King Charles Spaniel Muscular Dystrophy, (CKCS-MD)	X-linked Recessive	Clear
Centronuclear Myopathy, (CNM); mutation originally found in Great Dane	Autosomal Recessive	Clear
Centronuclear Myopathy, (CNM); mutation originally found in Labrador Retriever	Autosomal Recessive	Clear
Duchenne or Dystrophin Muscular Dystrophy, (DMD); mutation originally found in Golden Retriever	X-linked Recessive	Clear
Duchenne or Dystrophin Muscular Dystrophy, (DMD); mutation originally found in Norfolk Terrier	X-linked Recessive	Clear
Muscular Dystrophy, Ullrich-type; mutation originally found in Landseer	Autosomal Recessive	Clear
Myotonia Congenita; mutation originally found in Australian Cattle Dog	Autosomal Recessive	Clear
Myotonia Congenita; mutation originally found in Miniature Schnauzer	Autosomal Recessive	Clear
Myotubular Myopathy; mutation originally found in Rottweiler	X-linked Recessive	Clear
Nemaline Myopathy; mutation originally found in American Bulldog	Autosomal Recessive	Clear
X-Linked Myotubular Myopathy	X-linked Recessive	Clear



### Neurological Disorders - page 1

Disorder	Mode of Inheritance	Result	
Acral Mutilation Syndrome, (AMS)	Autosomal Recessive	Clear	
Alaskan Husky Encephalopathy, (AHE)	Autosomal Recessive	Clear	
Alexander Disease (AxD); mutation originally found in Labrador Retriever	Autosomal Dominant	Clear	
Bandera's Neonatal Ataxia, (BNAt)	Autosomal Recessive	Clear	
Benign Familial Juvenile Epilepsy or Remitting Focal Epilepsy	Autosomal Recessive	Clear	
Cerebellar Cortical Degeneration, (CCD); mutation originally found in Vizsla	Autosomal Recessive	Clear	
Cerebral Dysfunction; mutation originally found in Friesian Stabyhoun	Autosomal Recessive	nal Recessive Clear	
Dandy-Walker-Like Malformation (DWLM); mutation originally found in Eurasier	Autosomal Recessive	Clear	
Early-Onset Progressive Polyneuropathy; mutation originally found in Alaskan Malamute	Autosomal Recessive	Clear	
Fetal Onset Neuroaxonal Dystrophy, (FNAD)	Autosomal Recessive	Clear	
Hereditary Ataxia or Cerebellar Ataxia; mutation originally found in Old English Sheepdog and Gordon Setter	Autosomal Recessive	Clear	
Hyperekplexia or Startle Disease	Autosomal Recessive	Clear	
Hypomyelination; mutation originally found in Weimaraner	Autosomal Recessive	Clear	
Juvenile Myoclonic Epilepsy, (JME); mutation originally found in Rhodesian Ridgeback	Autosomal Recessive	Clear	
Juvenile encephalopathy; mutation originally found in Parson Russell Terrier	Autosomal Recessive	Clear	
L-2-Hydroxyglutaric aciduria, (L2HGA); mutation originally found in Staffordshire Bull Terrier	Autosomal Recessive	Clear	
L-2-Hydroxyglutaric aciduria, (L2HGA); mutation originally found in West Highland White Terrier	Autosomal Recessive	Clear	
Lagotto Storage Disease, (LSD)	Autosomal Recessive	Clear	
Neonatal Cerebellar Cortical Degeneration or Cerebellar Abiotrophy, (NCCD)	Autosomal Recessive	Clear	
Neonatal Encephalopathy with Seizures, (NEWS)	Autosomal Recessive	Clear	
Neuroaxonal Dystrophy (NAD); mutation originally found in Spanish Water Dog	Autosomal Recessive	Clear	



### Neurological Disorders - page 2

Disorder	Mode of Inheritance	Result
Neuronal Ceroid Lipofuscinosis 1, (NCL1); mutation originally found in Dachshund	Autosomal Recessive	Clear
Neuronal Ceroid Lipofuscinosis 10, (NCL10); mutation originally found in American Bulldog	Autosomal Recessive	Clear
Neuronal Ceroid Lipofuscinosis 5, (NCL5); mutation originally found in Border Collie	Autosomal Recessive	Clear
Neuronal Ceroid Lipofuscinosis 8, (NCL8); mutation originally found in Alpine Dachsbracke	Autosomal Recessive	Clear
Neuronal Ceroid Lipofuscinosis 8, (NCL8); mutation originally found in Australian Shepherd	Autosomal Recessive	Clear
Neuronal Ceroid Lipofuscinosis 8, (NCL8); mutation originally found in English Setter	Autosomal Recessive	Clear
Neuronal Ceroid Lipofuscinosis, (NCL7); mutation originally found in Chinese Crested Dog and Chihuahua	Autosomal Recessive	Clear
Polyneuropathy with ocular abnormalities and neuronal vacuolation, (POANV); mutation originally found in Black Russian Terrier	Autosomal Recessive	Clear
Progressive Early-Onset Cerebellar Ataxia; mutation originally found in Finnish Hound	Autosomal Recessive	Clear
Sensory Neuropathy; mutation originally found in Border Collie	Autosomal Recessive	Clear
Spinal Dysraphism	Autosomal Recessive	Clear
Spinocerebellar Ataxia with Myokymia and/or Seizures (SCA)	Autosomal Recessive	Clear
Spinocerebellar Ataxia/ Late-Onset Ataxia (SCA, LOA)	Autosomal Recessive	Clear
Spongy Degeneration with Cerebellar Ataxia, (SDCA1); mutation originally found in Belgian Shepherd Dog	Autosomal Recessive	Clear
X-Linked Tremors; mutation originally found in English Springer Spaniel	X-linked Recessive	Clear



#### **Neuromuscular Disorders**

Disorder	Mode of Inheritance	Result
Congenital Myasthenic Syndrome (CMS); mutation originally found in Labrador Retriever	Autosomal Recessive	Clear
Congenital Myasthenic Syndrome, (CMS); mutation originally found in Jack Russell Terrier	Autosomal Recessive	Clear
Congenital Myasthenic Syndrome, (CMS); mutation originally found in Old Danish Pointing Dog	Autosomal Recessive	Clear
GM1 Gangliosidosis; mutation originally found in Alaskan Husky	Autosomal Recessive	Clear
GM1 Gangliosidosis; mutation originally found in Portuguese Water Dog	Autosomal Recessive	Clear
GM1 Gangliosidosis; mutation originally found in Shiba Dog	Autosomal Recessive	Clear
GM2 Gangliosidosis, mutation originally found in Japanese Chin	Autosomal Recessive	Clear
GM2 Gangliosidosis; mutation originally found in Toy Poodle	Autosomal Recessive	Clear
Globoid Cell Leukodystrophy or Krabbe Disease, (GLD); mutation originally found in Irish Setter	Autosomal Recessive	Clear
Globoid Cell Leukodystrophy or Krabbe Disease, (GLD); mutation originally found in Terriers	Autosomal Recessive	Clear
Paroxysmal Dyskinesia, (PxD); mutation originally found in Irish Soft Coated Wheaten Terrier	Autosomal Recessive	Clear



#### **Skeletal Disorders**

Disorder	Mode of Inheritance	Result
Chondrodysplasia; mutation originally found in Norwegian Elkhound and Karelian Bear Dog	Autosomal Recessive	Clear
Cleft Palate; Cleft Lip and Palate with Syndactyly; ADAMTS20 gene mutation originally found in Nova Scotia Duck Tolling Retriever	Autosomal Recessive	Clear
Cleft Palate; DLX6 gene mutation originally found in Nova Scotia Duck Tolling Retriever	Autosomal Recessive	Clear
Craniomandibular Osteopathy, (CMO); mutation associated with terrier breeds	Autosomal Dominant (Incomplete Penetrance)	Clear
Hereditary Vitamin D-Resistant Rickets, (HVDRR)	Autosomal Recessive	Clear
Oculoskeletal Dysplasia 2 or Dwarfism-Retinal Dysplasia 2, (OSD2)	Autosomal Recessive	Clear
Osteochondrodysplasia; mutation originally found in Miniature Poodle	Autosomal Recessive	Clear
Osteochondromatosis; mutation originally found in American Staffordshire Terrier	Autosomal Dominant	Clear
Osteogenesis Imperfecta, (OI); mutation originally found in Beagle	Autosomal Dominant	Clear
Osteogenesis Imperfecta, (OI); mutation originally found in Dachshund	Autosomal Recessive	Clear
Skeletal Disease (Hypophosphatasia); mutation originally found in Karelian Bear Dog	Autosomal Recessive	Clear
Skeletal Dysplasia 2, (SD2)	Autosomal Recessive	Clear
Spondylocostal Dysostosis	Autosomal Recessive	Clear
Van den Ende-Gupta Syndrome, (VDEGS)	Autosomal Recessive	Clear



#### **Dermal Disorders**

Disorder	Mode of Inheritance	Result
Dystrophic Epidermolysis Bullosa; mutation originally found in Central Asian Ovcharka	Autosomal Recessive	Clear
Dystrophic Epidermolysis Bullosa; mutation originally found in Golden Retriever	Autosomal Recessive	Clear
Epidermolytic Hyperkeratosis	Autosomal Recessive	Clear
Focal Non-Epidermolytic Palmoplantar Keratoderma, (FNEPPK); mutation originally found in Dogue de Bordeaux	Autosomal Recessive	Clear
Hereditary Footpad Hyperkeratosis, (HFH)	Autosomal Recessive	Clear
Ichthyosis; mutation originally found in American Bulldog	Autosomal Recessive	Clear
Ichthyosis; mutation originally found in Great Dane	Autosomal Recessive	Clear
Lamellar Ichthyosis, (LI)	Autosomal Recessive	Clear
Ligneous Membranitis	Autosomal Recessive	Clear
Musladin-Lueke syndrome, (MLS)	Autosomal Recessive	Clear
X-Linked Ectodermal Dysplasia, (XHED)	X-linked Recessive	Clear



#### **Other Disorders**

Disorder	Mode of Inheritance	Result
Acute Respiratory Distress Syndrome, (ARDS); mutation originally found in Dalmatian	Autosomal Recessive	Clear
Amelogenesis Imperfecta, (AI); mutation originally found in Italian Greyhound	Autosomal Recessive	Clear
Amelogenesis Imperfecta, (AI); mutation originally found in Parson Russell Terrier	Autosomal Recessive	Clear
Congenital Keratoconjunctivitis Sicca and Ichthyosiform Dermatosis, (CKCSID)	Autosomal Recessive	Clear
Dental Hypomineralisation; mutation originally found in Border Collie	Autosomal Recessive	Clear
Narcolepsy; mutation originally found in Dachshund	Autosomal Recessive	Clear
Narcolepsy; mutation originally found in Doberman Pinscher	Autosomal Recessive	Clear
Narcolepsy; mutation originally found in Labrador Retriever	Autosomal Recessive	Clear
Persistent Müllerian Duct Syndrome, (PMDS); mutation originally found in Miniature Schnauzer	Autosomal Recessive	Clear
Primary Ciliary Dyskinesia, (PCD)	Autosomal Recessive	Clear



# APPENDIX Explanation of the results of the tested disorders

Autosomal recessive inheritance (ARI)

Clear - A dog carries no copies of the tested mutation and has no or reduced likelihood of developing and passing on the disease/condition.

Carrier - A dog carries one copy of the tested mutation. Carriers typically have a normal, healthy appearance but pass on the mutation to approximately 50% of their offspring.

At risk - A dog carries two copies of the tested mutation and is at high or increased risk of developing the disease/condition.

Autosomal dominant inheritance (ADI)

Clear - A dog carries no copies of the tested mutation and has no or reduced likelihood of developing and passing on the disease/condition.

At risk - A dog carries one or two copies of the tested mutation and is at high or increased risk of developing the disease/condition.

X-linked recessive inheritance (X-linked)

Clear - A dog carries no copies of the tested mutation and has no or reduced likelihood of developing and passing on the disease/condition.

Carrier - Female carriers typically have a normal, healthy appearance but carry one copy of the tested mutation on one of their X chromosomes. As males only have one X chromosome, there are no male carriers.

At risk - Female dogs at risk carry two mutated copies of the tested mutation. Males carry one copy of the tested mutation on their single X chromosome. Dogs at risk are at high or increased risk of developing the disease/condition.

Please note that the descriptions above are generalized based on typically observed inheritance patterns. When obtaining a 'carrier' or 'at risk' test result, always refer to the corresponding online test documentation for more detailed information on the condition and any exceptions.

### Genoscoper Laboratories - Legal Notice

Genoscoper Laboratories' services and test results are produced based on samples and materials supplied by the Client. Testing and analysis is performed by using methods and processes that Genoscoper Laboratories deems appropriate. Genoscoper Laboratories reserves the right to make changes in the collection of the single-gene tests included in the testing service as well as to remove results derived from them, if new information comes available that in any way questions the validity of the test results. Results provided by Genoscoper Laboratories are prepared solely for the use of the Client. For further information, please visit: <a href="https://www.mydogdna.com/legal-notices">www.mydogdna.com/legal-notices</a>

Registered Name: Sugarwhip Abydos

Nickname: Clyde Nickname: Clyde Registration ID: VDH/WCD 5608 Microchip: 276098106520463 DEU Breed: Whippet Gender: Male

Owner: Tina Sammt
Country: Germany
Testing date: 4/2/2018
DNA identification profile:
Identified with standard ISAG 2006 markers
Dog's identity verified from microchip or tattoo by veterinarian or other authorised person during sample taking: Yes





#### **Certificate of DNA Identification Profile**

17/8/2020

<b>REN64E19</b> 139/139	<b>REN247M23</b> 268/270	<b>REN169O18</b> 162/166	<b>REN169D01</b> 212/212	INU055 210/210	<b>INU030</b> 144/150	INU005 126/126		<b>AHT137</b> 137/147	<b>REN54P11</b> 236/236	<b>REN162C04</b> 206/206
REN105LO3 229/233	INRA21 99/101	<b>FH2054</b> 156/168	<b>CXX279</b> 116/118	,		<b>AHTH171</b> 227/227	<b>AHTH130</b> 127/131	<b>AHTK211</b> 87/87	<b>AHT121</b> 94/98	AMELOGENIN X/Y

On behalf of Genoscoper Laboratories,

Jonas Domeur SIGNATURE

Jonas Donner, PhD, Head of Research and Development at Genoscoper Laboratories

PROVIDED BY GENOSCOPER LABORATORIES PO Box. 1040 FI-00251 Helsinki | info@genoscoper.com